

1. A method for calibrating a biometric authentication device over time comprising the steps of:


Sub
Q1
obtaining an authenticating biometric value from an actual biometric measurement;
weighting the authenticating biometric value, and
5 integrating the weighted authenticating biometric value into an authenticating template.

2. The method of claim 1 wherein the step of obtaining an authenticated biometric value comprises determining that a measured biometric value falls within a predetermined range of biometric values.

3. The method of claim 2 wherein the integration of the weighted value is accomplished by averaging the weighted value into the authenticating range of values.

4. The method of claim 3 wherein the step of averaging the weighted value further comprises multiplying the authenticating measured value by a multiplier.

5. The method of claim 1 wherein the step of weighting the authenticating biometric value allows the biometric value to be adaptably weighted.



6. The method of claim 1 wherein biometric value is adaptably weighted based upon consistent differences in authenticated measured biometric values.

7. The method of claim 1 wherein biometric value is adaptably weighted based upon trends in measured authenticating biometric values.

8. The method of claim 1 wherein biometric value is adaptably weighted based on the frequency of use of the biometric authentication device.


9. The method of claim 1 wherein biometric value is adaptably weighted based on the number of uses of the biometric authentication device.

10. The method of claim 1 wherein the authenticated biometric values are obtained by measuring histological biometric markers.

11. The method of claim 1 wherein the authenticating biometric values are obtained measuring a physiological biometric marker.

12. The method of claim 1 wherein the authenticated biometric values are weighted to accommodate for known changes in a biometric marker.

13. The method of claim 1 wherein the biometric values are univariate values.




14. The method of claim 1 wherein the biometric values are multivariate values.

15. The method of claim 1 wherein the biometric marker being measured is an internal biometric marker.

5

1066007 "03231650



16. A device employing a biometric access system, said access system being adaptable to changes in a user's biometric over time, said device adapting to said changes by using the steps

of:

obtaining an authenticating biometric value from an actual biometric measurement;

weighting the authenticating biometric value,


integrating the weighted authenticating biometric value into an authenticating template.

17. The method of claim 16 wherein the step of obtaining an authenticated biometric value comprises determining that a measured biometric value falls within a predetermined range of biometric values.

18. The method of claim 17 wherein the integration of the weighted value is accomplished by averaging the weighted value into the authenticating range of values.

19. The method of claim 18 wherein the step of averaging the weighted value further comprises multiplying the authenticating measured value by a multiplier.

20. The method of claim 16 wherein the biometric marker being measured is an internal biometric marker.



21. The method of claim 16 wherein the step of weighting the authenticating biometric value allows the biometric value to be adaptably weighted.

22. The method of claim 16 wherein biometric value is adaptably weighted based upon
5 consistent differences in authenticated measured biometric values.


23. The method of claim 16 wherein biometric value is adaptably weighted based upon trends in measured authenticating biometric values.

24. The method of claim 16 wherein biometric value is adaptably weighted based on the frequency of use of the biometric authentication device.

25. The method of claim 16 wherein biometric value is adaptably weighted based on the number of uses of the biometric authentication device.

26. The method of claim 16 wherein the authenticated biometric values are obtained by measuring histological biometric markers.

27. The method of claim 16 wherein the authenticating biometric values are obtained
20 measuring a physiological biometric marker.




28. The method of claim 16 wherein the authenticated biometric values are weighted to accommodate for known changes in a biometric marker.

29. The method of claim 16 wherein the biometric values are univariate values.

5

30. The method of claim 16 wherein the biometric values are multivariate values.

2001 OCT 09 10 00 AM



36. The method of claim 31 wherein the step of weighting the authenticating biometric value allows the biometric value to be adaptably weighted.

37. The method of claim 31 wherein biometric value is adaptably weighted based upon
5 consistent differences in authenticated measured biometric values.

38. The method of claim 31 wherein biometric value is adaptably weighted based upon trends in measured authenticating biometric values.

10 39. The method of claim 31 wherein biometric value is adaptably weighted based on the frequency of use of the biometric authentication device.

15 40. The method of claim 31 wherein biometric value is adaptably weighted based on the number of uses of the biometric authentication device.

41. The method of claim 31 wherein the authenticated biometric values are obtained by measuring histological biometric markers.

20 42. The method of claim 31 wherein the authenticating biometric values are obtained measuring a physiological biometric marker.

